

A CHARACTERIZATION OF EXTREMAL SETS IN HILBERT SPACES

by Viet NguyenKhac ¹ & Khiem NguyenVan ²

Hanoi Institute of Mathematics & Hanoi University of Education

1991 Mathematics Subject Classification. Primary 46B20, 46E30.

Abstract

We give a characterization of extremal sets in Hilbert spaces that generalizes a classical theorem of H. W. E. Jung. We investigate also the behaviour of points near to the circumsphere of such a set with respect to the Kuratowski and Hausdorff measures of non-compactness.

1. Introduction

Let $(X, \|\cdot\|)$ be a Banach space. For a non-empty bounded subset A of X and a non-empty subset B of X we shall use the following notations: $d(A) := \sup \{\|x - y\| : x, y \in A\}$ – the diameter of A ; $r_B(A) := \inf_{y \in B} \sup_{x \in A} \|x - y\|$ – the relative Chebyshev radius of A with respect to B , in particular $r(A) := r_{\overline{\text{co}}A}(A)$ with $\overline{\text{co}}A$ denoting the closed convex hull of A ; $C_B(A) := \{y \in B : \sup_{x \in A} \|x - y\| = r_B(A)\}$ – the set of Chebyshev centers of A in B .

The Jung constant of X is defined by $J(X) := \sup \{r_X(A) : A \subset X, \text{ with } d(A) = 1\}$. In connection with uniform normal structure one considers also the following important geometric constant – the self-Jung constant of X (cf. [2]): $J_s(X) := \sup \{r(A) : A \subset X, \text{ with } d(A) = 1\}$. In the case X is an inner product space, it is known that $C_X(A)$ consists of a unique point which belongs to the closed convex hull $\overline{\text{co}}A$ of A (cf. [5]). Hence $J(X) = J_s(X)$ in this case. Classical Jung's theorem states that for $X = E^n$ – an n -dimensional Euclidean space we have $J(E^n) = J_s(E^n) = \sqrt{\frac{n}{2(n+1)}}$ ([7], cf. [4]). Furthermore if $X = H$ – a Hilbert space (the infinite-dimensional case), then $J(H) = J_s(H) = \frac{1}{\sqrt{2}}$ ([8], cf. [1], [2], [3]).

¹Institute of Mathematics, P.O.Box 631 Bo Ho, 10000 Hanoi, Vietnam; e-mail: nkviet@thevinh.ncst.ac.vn

²Department of Mathematics and Informatics, Hanoi University of Education, Cau Giay dist., Hanoi, Vietnam.

Definition. We say that a bounded subset A of X consisting of at least two points is an extremal set (resp. self-extremal set), if $r_X(A) = J(X)d(A)$ (resp. $r(A) = J_s(X)d(A)$).

Note that in the case $X = E^n$, or H , two these notions coincide, so in this case we shall speak simply of extremal sets. From the second part of the mentioned theorem of Jung one knows that a bounded subset A of E^n is extremal if and only if A contains a regular n -simplex with edges of length $d(A)$. In the case $X = H$ a partial result in this direction was obtained by Gulevich ([6]) who showed that if A is a relatively compact set in a Hilbert space with $d(A) > 0$, then $r(A) < \frac{1}{\sqrt{2}}d(A)$. Therefore if A is an extremal set in a Hilbert space, then A is not relatively compact.

Our aim in this paper is to give a characterization of extremal sets in Hilbert spaces which is a generalization of the second part of Jung's theorem.

Main Theorem. *Let A be an extremal set in a Hilbert space H with $d(A) = \sqrt{2}$. Then $\chi(A) = 1$ and for every $\varepsilon \in (0, \sqrt{2})$, for every positive integer p there exists a p -simplex Δ with its vertices in A and each edge of Δ has length not less than $\sqrt{2} - \varepsilon$.*

Conversely if $d(A) = \sqrt{2}$ and for every $\varepsilon \in (0, \sqrt{2})$, for every positive integer p there exists a p -simplex Δ with its vertices in A such that the length of each edge of Δ is not less than $\sqrt{2} - \varepsilon$, then A is an extremal set.

In the above formulation $\chi(A)$ denotes the Hausdorff measure of non-compactness of A , i.e. the infimum of positive r such that A can be covered by a finite number of balls with radius r and with centers in X . Besides, based on an observation of [3] ("Mushroom Lemma") we prove also a result on the behaviour of points near to the circumsphere of such a set with respect to the measure of non-compactness which says roughly that the main contribution to the measure of non-compactness comes from that part of the extremal set.

2. Measures of non-compactness of extremal sets

Theorem 1. *Let A be an extremal set in a Hilbert space H with $r(A) = 1$. Then we have $\alpha(A) = \sqrt{2}$.*

Here $\alpha(A)$ denotes the Kuratowski measure of non-compactness, that is the infimum of positive d such that A can be covered by a finite number of sets of diameter d .

The first proof. From the assumption $r(A) = 1$ it follows that for each integer number $n \geq 2$ we have $\bigcap_{x \in A} B(x, 1 - \frac{1}{n}) = \emptyset$, where $B(x, r)$ denotes the closed ball centered at x with radius r which is weakly compact since H is reflexive. Hence there exist $x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}$ in A such that $\bigcap_{i=p_{n-1}+1}^{p_n} B(x_i, 1 - \frac{1}{n}) = \emptyset$ (with convention $p_1 = 0$).

Setting $A_n := \{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}$ we denote the Chebyshev center of A_n in H by c_n and let $r_n := r(A_n)$, then $r_n > 1 - \frac{1}{n}$.

Let $S(c, r)$ denote the sphere with center c and radius r . From the proof of classical Jung's theorem one knows that $A \cap S(c_n, r_n) \neq \emptyset$ and $c_n \in \text{co}(A_n \cap S(c_n, r_n))$. So there exist $y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}$ in $A_n \cap S(c_n, r_n)$ (with convention $q_1 = 0$) and positive numbers $t_{q_{n-1}+1}, t_{q_{n-1}+2}, \dots, t_{q_n}$ such that

$$c_n = \sum_{q_{n-1} < i \leq q_n} t_i y_i, \quad \sum_{q_{n-1} < i \leq q_n} t_i = 1.$$

We claim that $\alpha(\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty) = \sqrt{2}$. Suppose on the contrary $\alpha(\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty) < \sqrt{2}$. Then one can choose $\varepsilon_0 \in (0, \sqrt{2})$ satisfying $\alpha(\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty) \leq \sqrt{2} - \varepsilon_0$, and so subsets $D_1, D_2, \dots, D_m$ of H with $d(D_i) \leq \sqrt{2} - \varepsilon_0$ for every $i = 1, 2, \dots, m$ such that $\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty \subset \bigcup_{i=1}^m D_i$. There exists at least one set among $D_1, D_2, \dots, D_m$, say D_1 with the property that there are infinitely many n satisfying

$$\sum_{i \in J_n} t_i \geq \frac{1}{m} \tag{1}$$

where

$$J_n := \{i \in [q_{n-1} + 1, q_n] : y_i \in D_1\}.$$

For each n satisfying (1) and fixed $j \in J_n$ we have

$$\begin{aligned} \sum_{q_{n-1} < i \leq q_n} t_i \|y_i - y_j\|^2 &= \sum_{q_{n-1} < i \leq q_n} t_i \|y_i - c_n + c_n - y_j\|^2 = \\ &= \sum_{q_{n-1} < i \leq q_n} t_i (\|y_i - c_n\|^2 + \|y_j - c_n\|^2 - 2(y_i - c_n, y_j - c_n)) = \\ &= 2r_n^2 - 2 \left(\sum_{q_{n-1} < i \leq q_n} t_i y_i - c_n, y_j - c_n \right) = \\ &= 2r_n^2 > 2 \left(1 - \frac{1}{n}\right)^2 > 2 - \frac{4}{n}, \end{aligned}$$

where $(\cdot, \cdot)$ denotes the inner product in H .

On the other hand

$$\begin{aligned}
\sum_{q_{n-1} < i \leq q_n} t_i \|y_i - y_j\|^2 &= \sum_{i \in J_n} t_i \|y_i - y_j\|^2 + \sum_{q_{n-1} < i \leq q_n, i \notin J_n} t_i \|y_i - y_j\|^2 \leq \\
&\leq (\sqrt{2} - \varepsilon_0)^2 \sum_{i \in J_n} t_i + 2 \left(1 - \sum_{i \in J_n} t_i\right) = \\
&= 2 - [2 - (2 - \varepsilon_0)^2] \left(\sum_{i \in J_n} t_i\right) \leq \\
&\leq 2 - [2 - (2 - \varepsilon_0)^2] \frac{1}{m}.
\end{aligned}$$

Hence

$$2 - [2 - (2 - \varepsilon_0)^2] \frac{1}{m} > 2 - \frac{4}{n}$$

with fixed ε_0 , m and for all n satisfying (1), a contradiction.

Thus $\alpha(\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty) = \sqrt{2}$. Since $d(A) = \sqrt{2}$ one concludes therefore $\alpha(A) = \sqrt{2}$. $\square$

For the second proof we need the following lemma which is a variation of [3, Lemma 4].

Lemma 2. *Let A be a non-empty bounded subset of Hilbert space H ; r and c the Chebyshev radius of A with respect to H and Chebyshev center of A in H , respectively. Then $c \in \overline{\text{co}}A_\varepsilon$ and $r = r(A_\varepsilon)$ for every $\varepsilon \in (0, r)$, where $A_\varepsilon := A \setminus B(c, r - \varepsilon)$.*

Proof of Lemma 2. Assume contrariwise that c is not the Chebyshev center of A_ε in H , then $r_1 := r(A_\varepsilon) < r$. Denoting by c_1 the Chebyshev center of A_ε in H we choose $c' = \alpha c_1 + (1 - \alpha)c$ for some $\alpha \in (0, 1)$ such that $0 < \|c - c'\| < \varepsilon$.

Take a point $x \in A$. If $x \in A_\varepsilon$ then $\|x - c'\| \leq \alpha \|x - c_1\| + (1 - \alpha) \|x - c\| \leq \alpha r_1 + (1 - \alpha)r < r$. In the other case $x \in A \setminus A_\varepsilon$ we have $\|x - c'\| \leq \|x - c\| + \|c - c'\| < r - \varepsilon + \|c - c'\| < r$.

So $A \subset B(c', r')$ with $r' < r$, a contradiction. The proof of the lemma is complete. $\square$

The second proof of Theorem 1. In view of Lemma 2 by taking $\varepsilon = \frac{1}{n}$ for every integer $n \geq 2$ one has: $c \in \overline{\text{co}}\left(A \setminus B\left(c, 1 - \frac{1}{n}\right)\right)$. Hence there exist $x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}$ in $A \setminus B\left(c, 1 - \frac{1}{n}\right)$ and positive numbers $t_{p_{n-1}+1}, t_{p_{n-1}+2}, \dots, t_{p_n}$ (with convention $p_1 = 0$) such that

$$\sum_{p_{n-1} < i \leq p_n} t_i = 1, \quad \left\| \sum_{p_{n-1} < i \leq p_n} t_i y_i - c \right\| < \frac{1}{n}.$$

We show that $\alpha(\{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}_{n=2}^\infty) = \sqrt{2}$. Assume on the contrary that $\alpha(\{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}_{n=2}^\infty) < \sqrt{2}$. By choosing $\varepsilon_0 \in (0, \sqrt{2})$ satisfying inequality $\alpha(\{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}_{n=2}^\infty) \leq \sqrt{2} - \varepsilon_0$, there exist subsets $D_1, D_2, \dots, D_m$ of H with $d(D_i) \leq \sqrt{2} - \varepsilon_0$ for every $i = 1, 2, \dots, m$, such that $\{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}_{n=2}^\infty \subset \bigcup_{i=1}^m D_i$. As in the first proof one can find among $D_1, D_2, \dots, D_m$, a set, say D_1 with the property that there are infinitely many n satisfying

$$\sum_{i \in I_n} t_i \geq \frac{1}{m} \quad (2)$$

where

$$I_n := \{i \in [p_{n-1} + 1, p_n] : x_i \in D_1\}.$$

Analogously for each n satisfying (2) and fixed $j \in I_n$ we have

$$\begin{aligned} \sum_{p_{n-1} < i \leq p_n} t_i \|x_i - x_j\|^2 &= \sum_{p_{n-1} < i \leq p_n} t_i \|y_i - c + c - y_j\|^2 = \\ &= \sum_{p_{n-1} < i \leq p_n} t_i (\|y_i - c\|^2 + \|y_j - c\|^2 - 2(y_i - c, y_j - c)) > \\ &> 2\left(1 - \frac{1}{n}\right)^2 - 2\left(\sum_{p_{n-1} < i \leq p_n} t_i x_i - c, y_j - c\right) \geq \\ &\geq \left(1 - \frac{1}{n}\right)^2 - 2\frac{1}{n} > 2 - \frac{6}{n}. \end{aligned}$$

Similarly one has also

$$\sum_{p_{n-1} < i \leq p_n} t_i \|x_i - x_j\|^2 \leq 2 - [2 - (\sqrt{2} - \varepsilon_0)^2] \frac{1}{m}$$

Hence

$$2 - [2 - (2 - \varepsilon_0)^2] \frac{1}{m} > 2 - \frac{6}{n}$$

for all n satisfying (2), a contradiction.

Thus $\alpha(\{x_{p_{n-1}+1}, x_{p_{n-1}+2}, \dots, x_{p_n}\}_{n=2}^\infty) = \sqrt{2}$. This implies $\alpha(A) = \sqrt{2}$. $\square$

As an immediate consequence one obtains Gulevich's result mentioned in the Introduction.

Corollary ([6]). *Let A be a relatively compact set in a Hilbert space with $d(A) > 0$. Then $r(A) < \frac{1}{\sqrt{2}}d(A)$.*

Remarks. 1. In [3] another proof of equality $J(H) = J_s(H) = \frac{1}{\sqrt{2}}$ was given by H. Steinlein. Essentially the heart of the proof is a relation between the Lifshitz characteristic and self-Jung constant of a Banach space X : $\varkappa_0(X) \leq (J_s(X))^{-1}$ which can be extended to the case of metric spaces with convex structure. We shall come back to this problem in a forthcoming paper.

2. By using Lemma 2 we see that A_ε is also an extremal set and $\alpha(A_\varepsilon) = \sqrt{2}$ for every $\varepsilon \in (0, 1)$.

3. Although $\alpha(A_\varepsilon) = \sqrt{2}$ for all $\varepsilon \in (0, 1)$, we may have $\overline{co}A \cap S(c, 1) = \emptyset$ (cf. [3]). The following question arises: what can be said about $\alpha(A \cap S(c, 1))$ when $A \cap S(c, 1) \neq \emptyset$? The answer is: $\alpha(A \cap S(c, 1))$ can take arbitrary values in $[0, \sqrt{2}]$. Below we produce some examples.

Example 1. Let $\{e_n\}_{n=1}^\infty$ be an infinite orthonormal sequence in Hilbert space H . Set $A_1 := \{(1 - \frac{1}{n})e_n\}_{n=1}^\infty$, and $A_2 := \{x_1, x_2, \dots, x_n, \dots\}$ with

$$\begin{aligned} x_1 &:= \frac{1}{\sqrt{2}}e_1 + \frac{1}{\sqrt{2}}e_2; & x_2 &:= \frac{1}{\sqrt{2}}e_1 + \frac{1}{2}e_2 + \frac{1}{2}e_3; \\ x_n &:= \frac{1}{\sqrt{2}}e_1 + \frac{1}{\sqrt{2^2}}e_2 + \dots + \frac{1}{\sqrt{2^n}}e_n + \frac{1}{\sqrt{2^n}}e_{n+1}; & \dots \end{aligned}$$

It is easy to see that $r(A_1) = 1$, $d(A_1) = \sqrt{2}$ and 0 is the Chebyshev center of A_1 in H . Furthermore $\|x_n\| = 1$ for every n , $\|x_m - (1 - \frac{1}{n})e_n\| \leq \sqrt{2}$, $\forall m, n$; $\|x_{n+p} - x_n\|^2 = \frac{1}{2^n} \rightarrow 0$ as $n \rightarrow \infty$. Thus $\{x_n\}$ is a Cauchy sequence and one gets $\alpha(A_2) = 0$.

Now setting $A := A_1 \cup A_2$ we have $r(A) = 1$, $d(A) = \sqrt{2}$ and 0 is also the Chebyshev center of A in H . Obviously $A \cap S(0, 1) = \overline{co}A \cap S(0, 1) = A_2$.

Example 2. Let $\{e_n\}_{n=1}^\infty$ and A_1 be as in Example 1. For each $\gamma \in (0, \sqrt{2}]$ putting $\beta := \frac{\gamma}{\sqrt{2}} \in (0, 1]$ we choose $\lambda \in [0, 1)$ satisfying $\lambda^2 + \beta^2 = 1$. Denote by $A_2 := \{y_1, y_2, \dots, y_n, \dots\}$ with

$$y_1 := \lambda e_1 + \beta e_2; \quad y_2 := \lambda e_1 + \beta e_3; \quad \dots \quad y_n := \lambda e_1 + \beta e_{n+1}; \quad \dots$$

Obviously $\|y_n\| = 1$ for every n ; $\|y_n - y_m\| = \sqrt{2}\beta = \gamma$, $\forall m \neq n$. Setting $A := A_1 \cup A_2$

we obtain $r(A) = 1$, $d(A) = \sqrt{2}$; 0 is the Chebyshev center of A in H and $A \cap S(0, 1) = \overline{c\partial}A \cap S(0, 1) = A_2$ with $\alpha(A_2) = \gamma$.

3. Proof of the Main Theorem

From the first proof of Theorem 1 we derived a sequence $y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}$ in $A_n \cap S(c_n, r_n)$ (with $q_1 = 0$) and positive $t_{q_{n-1}+1}, t_{q_{n-1}+2}, \dots, t_{q_n}$ for each integer $n \geq 2$ satisfying

$$c_n = \sum_{q_{n-1} < i \leq q_n} t_i y_i, \quad \sum_{q_{n-1} < i \leq q_n} t_i = 1.$$

We claim that $\chi(\{y_{q_{n-1}+1}, y_{q_{n-1}+2}, \dots, y_{q_n}\}_{n=2}^\infty) = 1$. Assume that A can be covered by a finite number of balls of radius r : $B_1, B_2, \dots, B_m$. Then there exist a ball among $B_1, B_2, \dots, B_m$, say B_1 , such that there are infinitely many n satisfying

$$\sum_{i \in J_n} t_i \geq \frac{1}{m} \quad (3)$$

where

$$J_n := \{i \in [q_{n-1} + 1, q_n] : y_i \in B_1\}.$$

As in the proof of Theorem 1 one has

$$\sum_{q_{n-1} < i \leq q_n} t_i \|y_i - y_j\|^2 = 2r_n^2 > 2 - \frac{4}{n} \quad (4)$$

for every $j \in [q_{n-1} + 1, q_n]$ fixed.

From (4) it follows that

$$\sum_{i \in I_{nj}} t_i < \frac{1}{\sqrt{n}},$$

where

$$I_{nj} := \left\{ i \in [q_{n-1} + 1, q_n] : \|y_i - y_j\|^2 < 2 - \frac{4}{\sqrt{n}} \right\}, \quad j \in [q_{n-1} + 1, q_n],$$

and

$$2(1 - t_j) \geq \sum_{q_{n-1} < i \leq q_n} t_i \|y_i - y_j\|^2 = 2r_n^2 > 2 - \frac{4}{n}.$$

This implies $t_j < \frac{2}{n}$ for every $j \in [q_{n-1} + 1, q_n]$. Therefore if n satisfies (3), then $|J_n| \frac{2}{n} > \frac{1}{m}$, or equivalently $|J_n| > \frac{n}{2m}$ (here $|J_n|$ denotes the cardinal of J_n).

For each n satisfying (3) and $j \in J_n$ let us denote by

$$J_n(y_j) := \left\{ i \in J_n : \|y_i - y_j\|^2 \geq 2 - \frac{4}{\sqrt{n}} \right\},$$

and

$$\hat{J}_n(y_j) := \{y_i : i \in J_n(y_j)\}.$$

Obviously from (5) one gets

$$\sum_{i \in J_n \setminus J_n(y_j)} t_i < \frac{1}{\sqrt{n}} \quad (6)$$

and

$$\sum_{i \in J_n(y_j)} t_i > \frac{1}{m} - \frac{1}{\sqrt{n}} \quad (7)$$

For each positive integer p choose n sufficiently large satisfying (3) and such that $\frac{p+1}{\sqrt{n}} \leq \frac{1}{\sqrt{m}}$. We claim that for every choice of $i_1, i_2, \dots, i_p \in J_n$ we have

$$\bigcap_{k=1}^p J_n(y_{i_k}) \neq \emptyset \quad (8)$$

Indeed otherwise $\bigcap_{k=1}^p J_n(y_{i_k}) = \emptyset$ would imply that

$$J_n(y_{i_1}) \subset J_n \setminus \left(\bigcap_{k=2}^p J_n(y_{i_k}) \right) = \bigcup_{k=2}^p (J_n \setminus J_n(y_{i_k})).$$

Consequently by (6) and (7)

$$\frac{1}{m} - \frac{1}{\sqrt{n}} < \sum_{\alpha \in J_n(y_{i_1})} t_\alpha \leq \sum_{k=2}^p \sum_{\alpha \in J_n \setminus J_n(y_{i_k})} t_\alpha < (p-1) \frac{1}{\sqrt{n}}.$$

Thus $\frac{1}{m} < \frac{p}{\sqrt{n}}$. This would contradict to the choice of n and p .

Next from (8) it follows that if $1 \leq k \leq p$ and $i_1, i_2, \dots, i_k \in J_n$, then $\bigcap_{\alpha=1}^k \hat{J}_n(y_{i_\alpha}) \neq \emptyset$. With n and p chosen as above let us fix $j \in J_n$. Setting $z_1 := y_j$ we take consecutively $z_2 \in \hat{J}_n(z_1)$; $z_3 \in \hat{J}_n(z_1) \cap \hat{J}_n(z_2)$; $\dots$; $z_{p+1} \in \bigcap_{i=1}^p \hat{J}_n(z_i)$.

Obviously $\|z_i - z_j\|^2 \geq 2 - \frac{4}{\sqrt{n}}$ for all $i \neq j$ in $\{1, 2, \dots, p+1\}$. Now for given $\varepsilon \in (0, \sqrt{2})$ choose n as above and moreover sufficiently large so that $2 - \frac{4}{\sqrt{n}} \geq (\sqrt{2} - \varepsilon)^2$. One sees that $z_1, z_2, \dots, z_{p+1}$ form a p -simplex Δ whose edges have length not less than $\sqrt{2} - \varepsilon$.

We now prove that the radius r of balls $B_1, B_2, \dots, B_m$ is ≥ 1 . Let c' and r' denote respectively the Chebyshev center of Δ in H and the Chebyshev radius of Δ with respect to H . From the proof of classical Jung's theorem it follows that there exist non-negative $\alpha_1, \alpha_2, \dots, \alpha_{p+1}$ with $\sum_{i=1}^{p+1} \alpha_i = 1$ and $c' = \sum_{i=1}^{p+1} \alpha_i z_i$. Next for each $j \in \{1, 2, \dots, p+1\}$ we have

$$\begin{aligned} \left(2 - \frac{4}{\sqrt{n}}\right)(1 - \alpha_j) &\leq \sum_{i=1}^{p+1} \alpha_i \|z_i - z_j\|^2 = \sum_{i=1}^{p+1} \alpha_i \|z_i - c' + c' - z_j\|^2 = \\ &= \sum_{i=1}^{p+1} \alpha_i (\|z_i - c'\|^2 + \|z_j - c'\|^2) - \left(\sum_{i=1}^{p+1} \alpha_i (z_i - c'), z_j - c' \right) \leq \\ &\leq 2(r')^2. \end{aligned}$$

Thus

$$\left(2 - \frac{4}{\sqrt{n}}\right) \sum_{j=1}^{p+1} (1 - \alpha_j) \leq 2(p+1)(r')^2,$$

or equivalently

$$r' \geq \sqrt{\frac{(2 - 4/\sqrt{n}) p}{2(p+1)}} \quad (9)$$

The RHS of (9) tends to 1 as $p \rightarrow \infty$. Obviously $r \geq r'$ since $\Delta \subset B_1$. This implies $r \geq 1$ as claimed. One concludes therefore $\chi(A) = 1$.

Conversly if $d(A) = \sqrt{2}$, and for every $\varepsilon \in (0, \sqrt{2})$ and every positive integer p A contains a p -simplex Δ with its edges having length $\geq \sqrt{2} - \varepsilon$, then we see immediately that A is an extremal set. $\square$

Acknowledgement. The second author would like to thank Professors Do Hong Tan and Nguyen Thanh Ha for encouragement and helpful discussions.

References

- [1] Berdyshev, V. I., *Connection between Jackson's inequality and a geometrical problem*, Math. Zametki, **3** (1968), 327–338 (Russian).
- [2] Bynum, W. L., *Normal structure coefficients for Banach spaces*, Pacific J. Math. , **86** (1980), 427–436.
- [3] Daneš, J., *On the radius of a set in a Hilbert space*, Comment. Math. Univ. Carolina, **25-2** (1984), 355–362.
- [4] Danzer, L., Grünbaum, B., Klee, V., *Helly's theorem and its relatives*, Proc. Symp. Pure Math., Amer. Math. Soc., **7** (1963), 101–180.
- [5] Garkavy, A. L., *On the Chebyshev center and convex hull of a set*, Uspekhi Math. Nauk, **19** (1964), N^o 6, 139–145.
- [6] Gulevich, N. M., *The radius of a compact set in a Hilbert space*, Zapiski Nauchnykh Seminarov LOMI im. V. A. Steklova, AN SSSR, vol. **164** (1988), 157–158 (Russian).
- [7] Jung, H. W. E., *Über die kleinste Kugel, die eine räumliche Figur einschliesst*, J. Reine Angew. Math., **123** (1901), 241–257.
- [8] Routledge, N. A., *A result in Hilbert space*, Quart. J. Math., Oxford, (2) **3** (1952), N^o 9, 12–18.